

- Cvjetković V, Wähner C, Lisgara M, Krejci R (2018c):** Comparison of lung lesions in slaughter pigs vaccinated with different *M. hyopneumoniae* vaccines during a yearly survey in Germany and Austria. Proceedings of the 25th IPVS 2018, Chongqing, China, 207.
- Dottori M, Nigrelli AD, Bonilauri P, Merialdi G, Gozio S, Cominotti F (2007):** Proposal for a new grading system for pleuritis at slaughterhouse. The S.P.E.S. (slaughterhouse Pleuritis evaluation system) grid. Large Anim Rev 13: 161–165.
- Eddicks M, Szikora F, Walhöfer N, Sauter Louis C, Reese S, Banholzer E, Reiner G, Sutter G, Ritzmann M, Fux R (2017):** Occurrence of genotypes of porcine circovirus (PCV2) in pig farms using different vaccination strategies against PCV2. Tierärztl Prax Ausg G Grosstiere Nutztiere 45: 90–97.
- Elicker S, Sipos W (2009):** The tissue compatibility of different *Mycoplasma hyopneumoniae* vaccines is mainly dependent upon their adjuvants. Berl Munch Tierärztl Wochenschr 122: 348–353.
- Elicker S, Scherer N, Sipos W (2009a):** Retrospektive Analyse der Ätiologie respiratorischer Erkrankungen österreichischer Absetzferkel. Tieraerztl Umsch 64: 484–489.
- Elicker S, Mayrhofer E, Scherer N, Fischer L, Weissenböck H, Sipos W (2009b):** Retrospektive Analyse der Ätiologie respiratorischer Erkrankungen von Mastschweinen sowie Jung- und Zuchtsauen aus Österreich. Wien Tierärztl Monat 96: 246–252.
- Fraille L, Sibila M, Nofrarias M, López-Jimenez R, Huerta E, Llorens A, López-Soria S, Pérez D, Segalés J (2012):** Effect of sow and piglet porcine circovirus type 2 (PCV2) vaccination on piglet mortality, viraemia, antibody titre and production parameters. Vet Microbiol 161: 229–234.
- Grosse Beilage E, Rohde N, Krieter J (2009):** Seroprevalence and risk factors associated with seropositivity in sows from 67 herds in north-west Germany infected with *Mycoplasma hyopneumoniae*. Prev Vet Med 88: 255–263.
- Hillen S, von Berg S, Köhler K, Reinacher M, Willems H, Reiner G (2014):** Occurrence and severity of lung lesions in slaughter pigs vaccinated against *Mycoplasma hyopneumoniae* with different strategies. Prev Vet Med 113: 580–588.
- Madec F, Kobisch M (1982):** Bilan lésionnel des poumons de porcs charcutiers à l'abattoir. Journées. Rech porcine en France 14: 405–412.
- Merialdi G, Dottori M, Bonilauri P, Luppi A, Gozio S, Pozzi P, Spaggiari B, Martelli P (2012):** Survey of pleuritis and pulmonary lesions in pigs at abattoir with a focus on the extent of the condition and herd risk factors. Vet J 193: 234–239.
- Ostanello F, Dottori M, Gusmara C, Leotti G, Sala V (2007):** Pneumonia disease assessment using a slaughterhouse lung-scoring method. J Vet Med A 54: 70–75.
- Reiner G, Bronnert B, Hohloch C, Fresen C, Haack I, Willems H, Reinacher M (2010):** Qualitative and quantitative distribution of PCV2 in wild boars and domestic pigs in Germany. Vet Microbiol 145: 1–8.

Korrespondenzadresse:

Prof. Dr. Wolfgang Sipos
 Klinisches Department für Nutztiere
 Veterinärmedizinische Universität Wien
 Veterinärplatz 1
 1210 Wien, Österreich
 wolfgang.sipos@vetmeduni.ac.at