



Malignant epithelioid mesothelioma in a Visayan spotted deer (*Cervus alfredi*)

*Malignes epitheloides Mesotheliom bei einem Prinz-Alfred-Hirsch (*Cervus alfredi*)*

Marine Anne Soléne Bazin*, Joshua Schwinn*, Bianca Kühl*, Federico Armando, Andreas Beineke

TABLE 1: Primary antibodies used for immunohistochemistry

Antibody	Clone	Dilution	Antigen retrieval	Source
Pan-Cytokeratin	Mouse, mAb (AE1/AE3)	1:500	P	Dako, Glostrup, Denmark
Cytokeratin 5/6	Mouse, mAb (D5/16B4)	1:100	MW	Dako, Glostrup, Denmark
Vimentin	Mouse, mAb (V9)	1:100	–	Dako, Glostrup, Denmark
Desmin	Mouse, mAb (D33)	1:100	–	Dako, Glostrup, Denmark
Smooth muscle actin	Mouse, mAb (1A4)	1:100	–	Dako, Glostrup, Denmark
Thyroid transcription factor-1	Mouse, mAb (8G7G3/1)	1:100	MW	Dako, Glostrup, Denmark
Calretinin	Rabbit, pAB	1:500	MW	Synaptic Systems, Germany

mAb: monoclonal antibody; pAb: polyclonal antibody; MW: pretreatment in citric buffer (pH: 6.0), heated for 20 minutes in a microwave oven; P: enzymatic pretreatment with pronase for 10 minutes